

ENHANCED PRODUCTION OF BARIUM IN LOW-MASS STARS: EVIDENCE FROM OPEN CLUSTERS

VALENTINA D'ORAZI

Dipartimento di Astronomia e Scienza dello Spazio, Università di Firenze, Firenze, Italy and
INAF –Osservatorio Astrofisico di Arcetri, Firenze, Italy

LAURA MAGRINI, SOFIA RANDICH, AND DANIELE GALLI

INAF –Osservatorio Astrofisico di Arcetri, Firenze, Italy

MAURIZIO BUSO

Dipartimento di Fisica, Università di Perugia, and Sezione INFN, Perugia, Italy

PAOLA SESTITO

INAF –Osservatorio Astrofisico di Arcetri, Firenze, Italy

Draft version January 20, 2009

ABSTRACT

We report the discovery of a trend of increasing barium abundance with decreasing age for a large sample of Galactic open clusters. The observed pattern of $[\text{Ba}/\text{Fe}]$ vs. age can be reproduced with a Galactic chemical evolution model only assuming a higher Ba yield from the s -process in low-mass stars than the average one suggested by parametrized models of neutron-capture nucleosynthesis. We show that this is possible in a scenario where the efficiency of the extra-mixing processes producing the neutron source ^{13}C is anti-correlated with the initial mass, with a larger efficiency for lower masses. This is similar to the known trend of extended mixing episodes acting in H-rich layers and might suggest a common physical mechanism.

Subject headings: Galaxy: disk, abundances, evolution; open clusters and associations: general, abundances

1. INTRODUCTION

Barium is one of the so-called heavy neutron-capture elements. It is formed by neutron captures occurring through two major mechanisms: (i) the slow capture process (s -process) and (ii) the rapid one (r -process). The r -process is thought to occur in high-mass stars, $M \gtrsim 8 M_{\odot}$, exploding as type-II SNe (e.g., Kratz et al. 2007; Travaglio et al. 1999). The s -process instead comes from low- and intermediate-mass stars, $1 M_{\odot} \lesssim M \lesssim 3 M_{\odot}$ (e.g., Busso et al. 2001), thanks to the repeated neutron expositions guaranteed by the activation of the neutron sources $^{13}\text{C}(\alpha, n)^{16}\text{O}$ and $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$. The s -process contribution from low-mass stars had a dominant role in the solar mixture of Ba nuclei, so that in the Sun Ba comes from the s -process at about the 80% level (Arlandini et al. 1999).

Barium abundances have been measured in stars with different metallicities (e.g., Burris et al. 2000 for halo stars; Cescutti et al. 2006 and Bensby et al. 2005 for thin and thick disk stars), constraining the nucleosynthetic processes at the origin of Ba. However, several details of the Ba production are still a matter of debate due to the free parameters affecting the models (see, e.g., for the r -process Mathews et al. 1992; Pagel & Tautvaisienė 1997; Travaglio et al. 1999; for the s -process Busso et al. 2001). In the s -process, the main source of uncertainty is the still unknown physical mechanism promoting the penetration of protons into the He inter-shell zone at dredge-up; this penetration is necessary for

explaining the formation of the main neutron source ^{13}C .

In this *Letter* we address the evolution of Ba with age by means of Ba measurements in a sample of open clusters (OCs) and by comparing the empirical pattern with the results of a Galactic chemical evolution model. We find an unexpected trend, in apparent conflict with the currently adopted average Ba yields. We also identify in which direction the average yields should be corrected and why this is physically justified.

2. SAMPLE AND ANALYSIS

Our sample (see Table 1) consists of 20 OCs spanning a wide interval in age (~ 30 Myr–8 Gyr), Galactocentric radius (7–22 kpc), and metallicity ($-0.3 \leq [\text{Fe}/\text{H}] \leq +0.4$). We analyzed a large collection of high-resolution spectra obtained by our group during the years with different instruments: UVES in slit-mode and FLAMES/UVES on the ESO VLT, CASPEC on the ESO 3.6m telescope, the Red-Long camera echelle spectrograph at CTIO, and SARG on the Italian National Telescope Galileo. The spectra of unevolved members were analyzed in the first 10 OCs in Table 1, while the analysis of the older and more distant clusters is based on giant stars.

Barium abundances were computed by means of equivalent width (EW) analysis and by using the driver *blends* in MOOG (Snedden, 1973–2002 version). The grid of 1-D LTE model atmospheres of Kurucz (1993) was employed. Our analysis uses two Ba lines: $\lambda\lambda$ 5853 and 6496 Å. As well known, Ba lines are affected by hyperfine-splitting (hfs); thus we included in our line list the hyperfine structure and isotopic splitting, following the approach

by McWilliam (1998) and using $\log gf$ values published there. An 81% s -process composition was adopted for the Sun and the sample stars. Stark and radiative broadening were treated in the standard way and for collisional damping we adopted the classical Unsöld (1955) approximation.

As a first step, we derived the solar Ba abundance using the same method as for the sample stars: for the Sun we adopted $T_{\text{eff}} = 5770$ K, $\log g = 4.44$ and $\xi = 1.1$ km/s (Randich et al. 2006), obtaining $\log n(\text{Ba}) = 2.22$ and $\log n(\text{Ba}) = 2.26$ for $\lambda = 5853$ and 6496 Å, respectively. The average solar value $\log n(\text{Ba}) = 2.24 \pm 0.02$ agrees very well with the meteoritic abundance of $\log n(\text{Ba}) = 2.22$ (Grevesse et al. 1996). Our analysis is strictly differential with respect to the Sun: we applied a line by line solar correction, canceling out the uncertainties on the adopted set of atomic parameters ($\log gf$). The feature at $\lambda = 6496$ Å always yields higher abundances, suggesting, as also stressed by Prochaska et al. (2000), the presence of an unidentified blend; however, the differential nature of our analysis should minimize this effect, in particular for dwarf stars with close-to-solar stellar parameters. Ba abundances were obtained adopting stellar parameters and uncertainties derived in our previous studies.

Errors in Ba abundances for each star are due to uncertainties in EWs and stellar parameters. The standard deviation from the mean of each cluster instead provides a good estimate of the global errors on the average abundances¹. In order to estimate systematic errors, we compared our results with measurements from the literature for six clusters. We found a good agreement within the uncertainties ($\Delta[\text{Ba}/\text{Fe}] < 0.06$ dex) for M67, Berkeley 20, and NGC 6253; the differences are instead larger for the Hyades ($\Delta[\text{Ba}/\text{Fe}] = -0.2$ dex), Collinder 261 ($\Delta[\text{Ba}/\text{Fe}] = +0.2$ dex), and Berkeley 29 ($\Delta[\text{Ba}/\text{Fe}] = +0.14$ dex). These results show that, whereas discrepancies with the literature are present, these are not systematic; in particular our Ba abundances are not systematically higher. Also, given the old age of Cr 261 and Be 29, our higher $[\text{Ba}/\text{Fe}]$ ratio for these clusters does not affect our final conclusions.

3. RESULTS

The results are summarized in Table 1, where we report the average $[\text{Ba}/\text{Fe}]$ ratios for each cluster along with the standard deviation from the mean. In Figure 1 we show $[\text{Ba}/\text{Fe}]$ ratios as a function of cluster age; different symbols indicate dwarf and giant members. In the figure we also show the solar $[\text{Ba}/\text{Fe}]$ as well as our own measurement for α Centauri performed by using EWs published by Porto de Mello et al. (Porto de Mello et al. 2008). Our Ba abundance is in excellent agreement with this previous result ($\Delta[\text{Ba}/\text{Fe}] < 0.02$ dex). The trend of increasing $[\text{Ba}/\text{Fe}]$ from the oldest clusters to the youngest ones is evident. The figure also shows that $[\text{Ba}/\text{Fe}]$ ratios are systematically higher for clusters where giants have been analyzed. We suggest that this might be due to the fact that our analysis is not strictly differential for giant stars, as their parameters are different from the solar




FIG. 1.— Average $[\text{Ba}/\text{Fe}]$ as function of age for the sample clusters; filled and open squares denote clusters whose analysis is based on dwarf and giant stars, respectively. We also show the solar value (solar symbol) and that of the α Centauri (star with errorbars).

ones.

First, the question arises whether the observed trend of increasing $[\text{Ba}/\text{Fe}]$ with decreasing age is real or rather due to any effect that, for a given intrinsic Ba abundance, would increase the observed EWs. In particular, members of the youngest clusters in our sample (ages below ~ 500 Myr) are characterized by strong magnetic fields and are chromospherically very active; thus Ba II line strengths might be affected by both a “magnetic intensification mechanism” (Landi degl’Innocenti & Landolfi 2004) and by non-LTE (NLTE) effects. In order to check whether magnetic intensification could be a problem, we measured the EWs of the weak Eu II 6437 Å feature in the sample stars. Magnetic intensification would cause a similar (if not larger) effect for Eu, whose hyper-fine structure is notoriously more complex (the number of Zeeman sub-levels is higher). We found that, at variance with Ba, Eu EWs do not increase with the age, thus excluding that this process affects Ba EWs.

On the other hand, NLTE effects might be important for Ba II features, in particular, for the $\lambda 6496$ Å line and for stars with close-to-solar metallicity (Mashonkina et al. 1999). It is very likely that NLTE effects are amplified in young stars due to the presence of hot chromospheres. While we cannot quantify this effect, we believe that $[\text{Ba}/\text{Fe}]$ for clusters younger than ~ 500 Myr are likely overestimated.

4. TIME EVOLUTION OF BA

In Figure 2 we plot the average $[\text{Ba}/\text{Fe}]$ of OCs (only those based on dwarf star analysis) and of field stars by Bensby et al. (2005) vs. stellar age. Although field stars show a larger scatter, the two patterns agree very well and our OCs do extend the same trend to younger ages. Qualitatively, the observed trend of Ba vs. age confirms

¹ The line list with the EWs measurements as well as stellar parameters, and Ba abundances of each single star along with errors are available upon request.




FIG. 2.— Average $[\text{Ba}/\text{Fe}]$ as function of stellar age for the subsample of clusters whose analysis is based on dwarfs (filled circles and inverted triangles) compared with the abundance pattern of disk stars (open triangles) by Bensby et al. (2005). Filled triangles represent abundance measurements that probably need NLTE corrections. The model results are shown for two set of yields: (a) standard yields (Travaglio et al. 1999; Busso et al. 2001), long-dashed curve; and (b) enhanced s -process yields, dot-dashed curve. Both models show a peak at old ages due to the r -process from massive stars.

the dominant role played, in Galactic disk stars, by the s -process occurring in low-mass red giants during their asymptotic giant branch (AGB) phase (Busso et al. 2001; Cescutti et al. 2006). For a more quantitative comparison, we also show in Fig. 2 the time evolution of $[\text{Ba}/\text{Fe}]$ computed with a Galactic chemical evolution model for the Milky Way. The model adopted is a generalization of the multi-phase model by Ferrini et al. (1992), originally built for the solar neighborhood, and later extended to model the entire Galaxy (Ferrini et al. 1994). We refer to Magrini et al. (2007, 2008) for a detailed description of the model. In our “standard” model we adopted the average s -process and r -process yields by Busso et al. (2001) (extrapolated as in Cescutti et al. 2006) and Travaglio et al. (2001), respectively. While the r -process yields do not affect the $[\text{Ba}/\text{Fe}]$ ratio in the metallicity range of our OCs (the r -process is important only for $[\text{Fe}/\text{H}] \lesssim -2.2$), the values of the s -process yields are crucial to reproduce the increasing trend of $[\text{Ba}/\text{Fe}]$ with decreasing age of the cluster.

The figure clearly shows that the model with standard yields does not reproduce the empirical distribution, and, in particular the observational evidence that at recent epochs Ba is sensitively enhanced both for field stars and for OCs. The standard s -process yields do not even fit the solar $[\text{Ba}/\text{Fe}]$ value and result in a Ba abundance ratio $[\text{Ba}/\text{Fe}] \approx -0.15$ at 1 Gyr, to be compared with the measured value of ~ 0.2 . In the original work from where the s -process yields were taken (Busso et al. 2001) it was underlined that at every epoch and for every group of stars there was evidence of a large spread in their efficiencies. However, a simply randomly distributed spread seems not to be the case: it would generate from each given stellar generation a yield close to the average one adopted by us, and this is clearly inadequate to explain the data. Instead, we “empirically” verified that the observations, showing a very late increase of

Ba in the Galaxy, would require an efficiency of production strongly weighted toward lower masses. Namely, the observations can be well fitted by increasing the yields of stellar masses between 1 and $1.5 M_{\odot}$ by a factor of ~ 6 with respect to the extrapolation obtained from the average yields (roughly corresponding to the case ST/1.5 in Busso et al. 2001).

The results of a chemical evolution model where we have adopted “enhanced s -process” yields are shown in Fig. 2. In this model the solar Ba abundance can be reproduced, with contributions from the s - and r -process of 84% and 16%, respectively. As mentioned, the model with standard yields has instead a lower percentage of Ba produced by the s -process at the time of formation of the Sun. More in general, the model with enhanced yields provides a recent production of Ba in better agreement with the observations of OCs older than ~ 500 Myr. Both models fail to fit younger clusters and, indeed, it would be difficult to imagine a process capable of creating Ba in the last 500 Myr of Galactic evolution, unless local enrichment is assumed. We suggest that the difference between our model predictions and the measured $[\text{Ba}/\text{Fe}]$ ratios for young clusters provides an estimate of the NLTE effects that affect the Ba determination in these stars.

We finally stress that the classical plots of $[\text{Ba}/\text{Fe}]$ against $[\text{Fe}/\text{H}]$ (see, e.g., Travaglio et al. 1999, 2001; Cescutti et al. 2006) suffer for a strong degeneration in terms of age, since they compress the most recent period. Conversely, the plot of $[\text{Ba}/\text{Fe}]$ against age favors the youngest ages and thus is ideal to approach our problem. We also investigated the presence of a possible relationship of $[\text{Ba}/\text{Fe}]$ with metallicity among our sample, finding that no significant trend is present between these two quantities.

5. DISCUSSION

Our results show for the first time there is a need for an enhanced Ba production in low-mass stars, mainly from 1 to $1.5 M_{\odot}$, formed from ~ 10 to ~ 5 Gyr ago (e.g. Charbonnel et al. 1996). We have shown that, when adopting the average yields that have been commonly used so far, Galactic evolutionary models are in sharp disagreement with data. An enhanced Ba production, like the one assumed in the previous section, seems to be necessary, but one has to verify whether this ad-hoc data-fitting is physically acceptable.

The most promising possibility to obtain enhancements in the Ba production, especially in low masses, is that the intrinsic spread found normally in the s -process efficiency at any metallicity be not really a spread, but rather a dependence of the efficiency itself on the initial mass, in the sense of higher production for lower masses. As time passes, lower mass stars with larger Ba production start to contribute, and the average yield changes in time. Specifically, the formation of the neutron source ^{13}C requires a penetration of protons below the formal convective border at dredge-up. This is a typical extra-mixing process, like those necessary to account for the carbon and oxygen isotopic mix in low-mass red giants, and for the production and destruction of Li in them. These mechanisms are known to be increasingly more efficient for decreasing stellar mass. The evidence coming from the Ba enhancement indirectly suggests that

the mixing processes driving the formation of the neutron source behave similarly, and are maybe originated by the same cause (e.g. Busso et al. 2007). In general, we verified that the efficiency in s -processing required in very low mass stars in order to explain the data is close to the upper limit (case ST*2) suggested by Busso et al (2001), not to the average of the spread there considered (case ST/1.5). This must come together with the (already known) increase in the dredge-up efficiency at low metallicity, so that this large amount of extra Ba can be carried to the surface and ejected into the ISM.

Other possibilities were also considered such as: i) an increase in the initial s -element abundances of the contributing stars, due to previous enrichment by mass transfer from more massive companions; and ii) an increase in the r -process seeds on which slow n -captures occur, assuming that old disk stars were still characterized by high r -process abundances. Both attempts proved incapable of increasing substantially the Ba yields.

6. CONCLUSIONS

High-resolution observations of OCs allowed us to derive Ba abundances in a uniform way in systems covering a wide age range. The $[\text{Ba}/\text{Fe}]$ derived in OCs (but also in field stars, by other authors) shows a steady increase with time. In the framework of Galactic chemical evolution, this can be explained by an enhanced s -process production of Ba in very old AGB stars below $1.5 M_{\odot}$.

This need can be reconciled with the present understanding of the s -process if an anti-correlation exists between the initial mass of the star and the effectiveness of the extra-mixing mechanism responsible for forming

the neutron source. Stars at the lowest mass limit for contributing would in this case have a higher efficiency in s -processing. Accounting for the observations then requires that this efficiency be close to the upper limit indicated by Busso et al. (2001), i.e. an amount of ^{13}C burnt per interpulse higher by a factor of three than the gross average (case ST/1.5) discussed in that work: this would produce an increase in the Ba yield by a factor of 5-6. This scenario is plausible if the mechanisms driving the formation of the ^{13}C pocket are similar to those promoting the slow circulations known to occur below the convective envelope.

The youngest clusters of our sample, with ages < 500 Myr, have even higher $[\text{Ba}/\text{Fe}]$ ratios. We suggest that this is due to NLTE effects, and, in particular, to the effects of a strong, warm flux from a chromosphere. The comparison with our model suggests that NLTE effects in young stars might increase the $[\text{Ba}/\text{Fe}]$ ratios by a factor of two or more. Independent estimates of these effects from NLTE calculations as well as a more complete analysis of the hypothesis here advanced of an anticorrelation between initial mass and s -process efficiency are strongly needed.

We are grateful to A. McWilliam and C. Sneden for many helpful suggestions. We warmly thank L. Belluzzi and G. Cauzzi for useful discussions. The anonymous referee is acknowledged for a careful reading of the paper and valuable comments. This work has made use of the WEBDA database, originally developed by J. C. Mermilliod, and now maintained by E. Paunzen.

REFERENCES

- Arlandini, C., Käppeler, F., Wisshak, K. 1999, *ApJ*, 525, 886
 Bensby, T., Feltzing, S., Lundström, I., Ilyin, I. 2005, *A&A*, 433, 185
 Bragaglia A., Sestito P., Villanova, S., Carretta, E., Randich, S., Tosi, M. 2008, *A&A*, 480, 79
 Busso, M., Gallino, R., Lambert, D.L., Travaglio, C., & Smith, V.V. 2001, *ApJ*, 557, 802
 Busso, M., Wasserburg, G.J., Nollet, K.M., Calandra, A. 2007, *ApJ*, 671, 802
 Burris, D. L., Pilachowski, C. A., et al. 2000 *ApJ*, 544, 302
 Cescutti, G., François, P., Matteucci, F., Cayrel, R., Spite, M. 2006, *A&A*, 448, 557
 Charbonnel, C., Meynet, G., Maeder, A., Schaerer, D. 1996, *A&AS*, 115, 339
 D’Orazi, V., Randich, S. 2008, *A&A*, submitted
 Ferrini, F., Matteucci, F., Pardi, C., Penco, U. 1992, *ApJ*, 387, 138
 Ferrini, F., Mollà, M., Pardi, M. C., Diaz, A. I. 1994, *ApJ*, 427, 745
 Grevesse, N., Noels, A., Sauval, A. J. 1996, *ASP Conf. Ser.* 99, 117
 Kratz, K.-L., Farouqi, K., Mashonkina, L.I., and Pfeiffer, B. 2007, *New Astr. Rev.* 52, 390
 Kurucz, R.L., 1993, CD-ROM Nr. 9
 Landi degl’Innocenti, E., Landolfi, M. 2004, *Polarization in spectral lines*, Kluwer Academic Publishers
 Magrini, L., Corbelli, E., Galli, D. 2007, *A&A*, 470, 843
 Magrini, L., Sestito, P., Randich, S., Galli, D. 2008, *A&A*, in press (arXiv:0812.0854)
 Mashonkina, L., Gehren, T., Bikmaev, I. 1999, *A&A*, 343, 519
 Mathews, G.J., Bazan, G., Cowan, J.J. 1992, *ApJ*, 391, 719
 McWilliam, A., 1998, *AJ*, 115, 1640
 Pace, G., Pasquini, L., François, P. 2008, *A&A*, 489, 403
 Pagel, B.E.J., Tautvaišienė, G. 1997, *MNRAS*, 288, 108
 Paulson, D.B., Sneden, C., Cochran, W.D. 2003, *AJ*, 125, 3185
 Porto de Mello, G.F., Lyra, W., Keller, G.R. 2008, *A&A*, 488, 653
 Prochaska, J.X., Naumov, S.O., Carney, B.W., McWilliam, A., Wolfe, A.M. 2000, *AJ*, 120, 2513
 Randich, S., Sestito, P., Primas, F., Pallavicini, R., Pasquini, L. 2006, *A&A*, 450, 557
 Randich, S., Primas, F., Pasquini, L., Sestito, P., Pallavicini, R. 2007, *A&A*, 469, 163
 Sestito, P., Randich, S., Mermilliod, J.C., Pallavicini, R. 2003, *A&A*, 407, 289
 Sestito, P., Randich, S., Pallavicini, R. 2004, *A&A*, 426, 809
 Sestito, P., Randich, S. 2005, *A&A*, 442, 615
 Sestito, P., Bragaglia, A., Randich, S., Carretta, E., Prisinzano, L., Tosi, M. 2006, *A&A*, 458, 121
 Sestito, P., Randich, S., Bragaglia, A. 2007, *A&A*, 465, 185S
 Sneden, C.A. 1973, *ApJ*, 184, 839
 Terndrup, D.M., Pinsonneault, M., Jeffries, R.D., et al. 2002, *ApJ*, 576, 950
 Travaglio, C., Galli, D., Gallino, R., et al. 1999, *ApJ*, 521, 691
 Travaglio, C., Galli, D., Burkert, A. 2001, *ApJ*, 547, 217
 Unsöld, A., 1955, *Physik der Sternatmosphären* (Berlin: Springer-Verlag)

TABLE 1
OC SAMPLE

Cluster	Age (Gyr)	[Fe/H]	Ref.	[Ba/Fe]	# stars	T_{eff} (K)
IC 2602	0.035	0 ± 0.01	1	0.64 ± 0.07	8	4770–5760
IC 2391	0.053	-0.01 ± 0.02	1	0.68 ± 0.07	6	4680–5970
NGC 2516	0.110	$0.01 \pm 0.07^{\text{a}}$	2	0.41 ± 0.04	4	5110–5659
NGC 6475	0.22	0.14 ± 0.06	3	0.33 ± 0.11	19	5048–5888
Hyades	0.7	$0.13 \pm 0.05^{\text{b}}$	2	0.30 ± 0.05	4	5079–5339
Praesepe	0.7	0.27 ± 0.04	4	0.22 ± 0.06	6	5720–6280
NGC 3680	1.4	-0.04 ± 0.03	4	0.25 ± 0.06	2	6010–6210
IC 4651	1.7	0.12 ± 0.05	4	0.20 ± 0.03	5	5910–6320
NGC 752	2.0	0.01 ± 0.04	5	0.19 ± 0.08	9	5531–6151
M67	4.3	0.02 ± 0.14	6	0.04 ± 0.05	10	5541–6223
NGC 2324	0.7	-0.17 ± 0.05	7	0.66 ± 0.09	7	4300–5100
NGC 3960	0.7	0.02 ± 0.04	8	0.50 ± 0.06	6	4870–5050
NGC 2660	0.8	0.04 ± 0.04	8	0.47 ± 0.06	5	5030–5200
NGC 2477	1.0	0.07 ± 0.00	7	0.40 ± 0.03	6	4950–5030
Berkeley 20	4.3	-0.30 ± 0.02	10	0.09 ± 0.20	2	4400–4770
Berkeley 29	4.3	-0.31 ± 0.03	10	0.40 ± 0.10	5	4970–5020
NGC 6253	4.5	0.39 ± 0.08	9	0.14 ± 0.10	5	4450–6200
Melotte 66	5.4	-0.33 ± 0.03	10	0.32 ± 0.10	6	4717–4850
Berkeley 32	6.1	-0.29 ± 0.04	8	0.24 ± 0.15	9	4760–4920
Collinder 261	8.4	0.13 ± 0.05	10	0.22 ± 0.03	7	4350–4720

NOTE. — Results of [Ba/Fe] ratios for our sample OCs. We list: ages (Col. 2) from Magrini et al. (2008), with the exception of NGC 2516 (WEBDA) and NGC 752 (Sestito et al. 2004); [Fe/H] values along with references for them and stellar parameters (Cols. 3 and 4); average [Ba/Fe] ratios along with the standard deviation (Col. 5); number of stars and range of effective temperatures (Cols. 6 and 7). Reference codes are as follows: 1=D’Orazi & Randich (2008), 2=Randich et al. (2007), 3=Sestito et al. (2003), 4=Pace et al. (2008), 5=Sestito et al. (2004), 6=Randich et al. (2006), 7=Bragaglia et al. (2008), 8=Sestito et al. (2006), 9=Sestito et al. (2007), 10=Sestito et al. (2008).